

SM Predictions for Muon $(g - 2)/2$

Simon Eidelman

Budker Institute of Nuclear Physics,
Novosibirsk, Russia

Outline

1. QED
2. Electroweak term
3. Progress with e^+e^- data
4. LO hadronic term
5. Prospects
6. Errors or New Physics?
7. Conclusions

Muon Anomalous Magnetic Moment

$$\vec{\mu} = g \frac{e}{2m} \vec{S}, \quad a = (g - 2)/2.$$

In Dirac theory for pointlike particles $g = 2$,
higher-order effects or new physics $\Rightarrow g \neq 2$

Any significant difference of a_{μ}^{exp} from a_{μ}^{th} indicates
New Physics beyond the Standard Model.

a_{μ} is much more sensitive to new physics effects than a_e :
the gain is usually $\sim (m_{\mu}/m_e)^2 \approx 4.3 \cdot 10^4$.

$$a_{\mu}^{\text{th}} = a_{\mu}^{\text{SM}} + a_{\mu}^{\text{NP}}, \quad a_{\mu}^{\text{SM}} = a_{\mu}^{\text{QED}} + a_{\mu}^{\text{EW}} + a_{\mu}^{\text{had}}.$$

Experimental Status of a_l

$$a_e = 1159652180.73(28) \times 10^{-12} \quad 0.24\text{ppt}$$

D. Hanneke et al., PRL 100, 120801 (2008)
QED test or α determination

$$a_\mu = 116592080(63) \times 10^{-11} \quad 0.54\text{ppb}$$

G.W. Bennett et al. (E821), PRD 73, 072003 (2006)
Sensitive test of the Standard Model

$$a_\tau = -0.018(17) \quad \text{or} \quad -0.052 < a_\tau < 0.013 \quad 95\%CL$$

J. Abdallah et al. (DELPHI), EPJ C 35, 159 (2004)
Theory: $117721(5) \times 10^{-8}$, SE, M. Passera, MPL A 22, 159 (2007)

QED Contribution a_μ^{QED}

$$\begin{aligned}
 a_\mu^{\text{QED}} \cdot 10^{10} = \sum C_i \left(\frac{\alpha}{\pi}\right)^i = & \quad 11614097.3 \text{ (1-loop)} & \quad 1 \text{ diagram} \\
 & + \quad 41321.8 \text{ (2-loop)} & \quad 9 \\
 & + \quad 3014.2 \text{ (3-loop)} & \quad > 100 \\
 & + \quad 38.1 \text{ (4-loop)} & \quad > 1000 \\
 & + \quad 0.4 \text{ (5-loop)} & \quad > 20000
 \end{aligned}$$

α^3 terms known analytically (S. Laporta, E. Remiddi, 1993),

α^4 terms – numerically (T. Kinoshita et al., 2003-2008),

$L \log \alpha^5$ (TK et al., 2005,2007; A.L. Kataev, 2006, K. Chetyrkin et al., 2008):


$$a_\mu^{\text{QED}} = (116584719.4 \pm 1.4) \cdot 10^{-11}.$$

From the latest value of a_e (D. Hanneke et al., 2008; M. Passera, 2008):

$$\alpha^{-1} = 137.035999084(51), \quad a_\mu^{\text{QED}} = (116584718.09 \pm 0.14 \pm 0.04) \cdot 10^{-11}.$$

The errors are due to: a/ $\mathcal{O}(\alpha^5)$, b/ α

Electroweak contribution a_μ^{EW}




One-loop electroweak contributions

Authors	Year	$a_\mu^{\text{EW}}, 10^{-10}$
..., ..., ...	1972	19.5
A. Czarnecki et al.	1996	15.2 ± 0.4
A. Czarnecki et al.	2002	$15.4 \pm 0.1 \pm 0.2$

The errors are due to: a/ hadr. loops, b/ M_H, M_t , 3-loop effects.

Hadronic contribution a_μ^{had}

$$a_\mu^{\text{had}} = a_\mu^{\text{had,LO}} + a_\mu^{\text{had,HO}} + a_\mu^{\text{had,LBL}}$$




$$a_\mu^{\text{had,LO}} = \left(\frac{\alpha m_\mu}{3\pi}\right)^2 \int_{4m_\pi^2}^{\infty} ds \frac{R(s) \hat{K}(s)}{s^2},$$

C. Bouchiat, L. Michel, Bouchiat, 1961;
M. Gourdin, E. de Rafael, 1969

$$R(s) = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)},$$

$\hat{K}(s)$ grows from 0.63 at $s = 4m_\pi^2$ to 1 at $s \rightarrow \infty$,
 $1/s^2$ emphasizes the role of low energies,
 particularly important is the reaction $e^+e^- \rightarrow \pi^+\pi^-$
 with a large cross section below 1 GeV.

Contributions of Various Energy Ranges to $a_{\mu}^{\text{had,LO}}$




How is R(s) Measured?

- $\sqrt{s} < 2$ GeV – exclusive modes
($\pi^+\pi^-$, $\pi^+\pi^-\pi^0$, ..., $K\bar{K}$, ...)
- Possibly missing (small σ , undetected) final states
- Above 2 GeV – total R
(all multihadronic events)
- Initial state radiation (ISR), vacuum polarization (VP), final state radiation (FSR):
M. Drees, K. Hikasa, 1990


$$\sigma_{\text{dr}} = \frac{N}{\int L dt \epsilon (1 + \delta_{\text{ISR}})}$$

$$\sigma_{\text{bare}} = \sigma_{\text{dr}} |1 - \Pi(s)|^2 (1 + \delta_{\text{FSR}})$$




$$|1 - \Pi(s)|^2$$

$e^+e^- \rightarrow \pi^+\pi^-$ (CMD-2, SND and KLOE)




CMD-2: $\sim 9 \cdot 10^5$ ev.

2E, MeV	$\sigma, \%$
370-520	0.7
600-970	0.6-0.8
1040-1380	1.3-4.2



SND: $\sim 8 \cdot 10^5$ ev.


2E, MeV	$\sigma, \%$
390-420	3.2
430-970	1.3





KLOE: $\sim 3.1 \cdot 10^6$ ev.
(590-970) MeV - 0.9%

Expecting news from KLOE (G.Venanzoni) and BaBar (M.Davier)

Hadronic Cross Sections at CMD-2






Most of these channels have also been studied
at SND with consistent results





Good agreement with SND. BaBar's points much higher than at DM2

Source	Before BaBar	All + BaBar	All-DM2+BaBar
$\delta a_\mu, 10^{-10}$	$2.45 \pm 0.26 \pm 0.03$	$2.79 \pm 0.19 \pm 0.01$	$3.25 \pm 0.09 \pm 0.01$



Good agreement with CMD-2/SND.
Error of continuum below 2 GeV improved.


$$e^+e^- \rightarrow 6\pi$$



δa_μ changed from 0.10 ± 0.10 to 0.108 ± 0.016 and from 1.42 ± 0.30 to 0.890 ± 0.093

Significant improvement compared to the previous data!

R Measurements at $\sqrt{s} < 10$ GeV



Excellent agreement with the pQCD predictions!

e^+e^- Data Based Calculation of $a_\mu^{\text{had,LO}}$ (DEHZ-2006)


\sqrt{s} , GeV	$a_\mu^{\text{had,LO}}, 10^{-10}$	$\delta a_\mu^{\text{had,LO}}, \%$
2π	$504.6 \pm 3.1 \pm 1.0$	73.0
ω	$38.0 \pm 1.0 \pm 0.3$	5.5
ϕ	$35.7 \pm 0.8 \pm 0.2$	5.2
0.6 – 1.8	$54.2 \pm 1.9 \pm 0.4$	7.8
1.8 – 5.0	$41.1 \pm 0.6 \pm 0.0$	6.0
$J/\psi, \psi'$	$7.4 \pm 0.4 \pm 0.0$	1.1
> 5.0	$9.9 \pm 0.2 \pm 0.0$	1.4
Total	$690.9 \pm 3.9_{\text{exp}} \pm 1.9_{\text{rad}} \pm 0.7_{\text{QCD}}$	100.0

Higher accuracy of e^+e^- data: the $a_\mu^{\text{had,LO}}$ error is 4.4 (0.63%) compared to 15.3 of EJ, 1995 and 7.2 of DEHZ, 2003!

Various $a_{\mu}^{\text{had,LO}}$ Calculations

Calculation	$a_{\mu}^{\text{had,LO}}, 10^{-10}$	$\sqrt{s}_{\text{pQCD}}, \text{ GeV}$
EJ, 1995	702.3 ± 15.3	40
DEHZ, 2003	696.3 ± 7.2	5
DEHZ, 2006	690.9 ± 4.4	5
HMNT, 2008	689.4 ± 4.6	11.09
J, 2008	692.3 ± 6.0	12


Higher Order Hadronic Contributions $a_{\mu}^{\text{had,HO}}$



The contributions of all 3 graphs can be calculated in terms of the $\int R(s)G(s)ds/s^{2(3)}$, where $G(s)$ is a smooth function of s , so that the low energy range again dominates the integral. Several calculations agree. The accepted value is (B. Krause, 1997; K. Hagiwara et al., 2003):

$$a_{\mu}^{\text{had,HO}} = (-9.8 \pm 0.1) \cdot 10^{-10}.$$

Light-by-Light Scattering



Various approaches used:

- Vector Dominance and Chiral models
- Data on $\gamma\gamma^* \rightarrow \pi^0, \eta, \eta'$ (single-tag)
- Effective field theory

Authors	Year	$a_\mu^{\text{lbl}}, 10^{-10}$
J. Bijnens et al.	1996 (2002)	8.3 ± 3.2
M. Hayakawa and T. Kinoshita	1998 (2002)	9.0 ± 1.5
K. Melnikov and A. Vainshtein	2003	13.6 ± 2.5
M. Davier and W. Marciano	2004	12.0 ± 3.5
J. Bijnens and J. Prades	2006	11.0 ± 4.0
F. Jegerlehner and A. Nyffeler	2008	11.9 ± 4.0

Theory vs Experiment – I


Contribution	$a_\mu, 10^{-10}$
Experiment	11659208.0 ± 6.3
QED	11658471.8 ± 0.016
Electroweak	$15.4 \pm 0.1 \pm 0.2$
Hadronic	693.1 ± 5.6
Theory	11659180.3 ± 5.6
Exp.–Theory	$27.7 \pm 8.4 (3.3\sigma)$

The difference between experiment and theory is 3.3σ !

Theory vs Experiment – II


Calculation	$a_{\mu}^{\text{exp}}, 10^{-10}$	$a_{\mu}^{\text{th}}, 10^{-10}$	$a_{\mu}^{\text{exp}} - a_{\mu}^{\text{th}}, 10^{-10} (\sigma)$
EJ, 1995	11659230 ± 84	11659210 ± 16	$20 \pm 86 (---)$
DEHZ, 2003	11659203 ± 8	11659180.9 ± 8.0	$22.1 \pm 11.3 (1.9)$
DEHZ, 2006	11659208.0 ± 6.3	11659180.3 ± 5.6	$27.7 \pm 8.4 (3.3)$
HMNT, 2008	11659208.0 ± 6.3	11659180.4 ± 5.1	$27.6 \pm 8.1 (3.4)$
J, 2008	11659208.0 ± 6.3	11659181.3 ± 7.2	$26.7 \pm 9.6 (2.8)$

Theory vs Experiment – III




How Real Is a_μ^{had} Accuracy?

- Missing states: neutrals;
 $\pi^+\pi^-\pi^0, K\bar{K}n\pi$ - isospin
- New states from BaBar, double counting
- Radiative corrections (FSR):
Charge asymmetry at KLOE
3k $e^+e^- \rightarrow \pi^+\pi^-\gamma$ evts at CMD-2
- Correlations
- Averaging
- Light-by-light term
- Double counting (LO and HO)




Charge asymmetry in $e^+e^- \rightarrow \pi^+\pi^-\gamma$
sQED is OK ($0.6 < M_{\pi\pi}^2 < 0.7 \text{ GeV}^2$)

CVC. $e^+e^- \rightarrow X^0$ and $\tau^- \rightarrow \nu_\tau X^-$



Allowed $I^G J^P = 1^+ 1^-$:
 $X^- = \pi^- \pi^0, (4\pi)^-, \omega \pi^-,$
 $\eta \pi^- \pi^0, K^- K^0, (6\pi)^-, \dots$



Large SU(2) breaking corrections from theory, V.Cirigliano et al., 2002

$M(\Gamma)_{\rho^0} \neq M(\Gamma)_{\rho^\pm}$ helps,


M.Davier, 2003; S.Ghozzi, F.Jegerlehner, 2004

Corrections to the τ Spectral Functions

- $S_{EW} = 1.0233 \pm 0.0006$
- Real photons, loops
- FSR
- $m_{\pi^\pm} \neq m_{\pi^0}$
(phase space, Γ_ρ)
- $m_{\rho^\pm} \neq m_{\rho^0}$
- $\rho - \omega$ interference
- Radiative decays
($\pi\pi\gamma, \pi(\eta)\gamma, l^+l^-$)
- $m_u \neq m_d$
and 2 class currents


V. Cirigliano, G. Ecker,
H. Neufeld, 2002

M. Davier, S. Eidelman,
A. Höcker, Z. Zhang, 2002



G.Lopez Castro, 2006

CVC in the 2π Channel. e^+e^- vs. τ (Branchings)





The branching from all groups is systematically higher than the CVC prediction:
 $\mathcal{B}_\tau - \mathcal{B}_{ee} = (0.94 \pm 0.21)\%$.
 The discrepancy – a 3.6% effect, twice the SU(2) correction.

The puzzle remains unsolved \Rightarrow τ data not used

M. Benayoun, EPJ C55, 199 (2008):
 more complete SU(2) breaking, mixing, no conflict

New data on $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ from Belle



$$\mathcal{B}_{\text{Belle}} = (25.24 \pm 0.04 \pm 0.39)\%$$

$$\mathcal{B}_{\text{ALEPH}} = (25.471 \pm 0.097 \pm 0.085)\%$$

The contributions to a_μ^{had} are also compatible due to compensation at tails

Future of $a_{\mu}^{\text{had}} - I$

What can be done from the e^+e^- side?

More ISR analysis from KLOE, BaBar, Belle; better R below 4.3 GeV from CLEO-c: 4.4 \rightarrow 2.8

Experiments at VEPP-2000 with 2 detectors up to $\sqrt{s}=2$ GeV with $L_{\text{max}} = 10^{32} \text{ cm}^{-2}\text{s}^{-1}$, $10^{30} \text{ cm}^{-2}\text{s}^{-1}$ achieved this June

A similar machine (DAΦNE-II) is discussed in Frascati, a $\tau - c$ factory in Beijing commissioning.

By 2012: 2.8 \rightarrow 2.2, the total error of 4.6 limited by the LBL term (4.0)

Future of $(g_\mu - 2)/2$ – II

1. New $(g_\mu - 2)/2$ experiment at FLAB expects the accuracy of $0.1\text{ppm}_{\text{stat}}$ and $0.1\text{ppm}_{\text{syst}}$ or 1.5×10^{-10}
2. Such accuracy for $a^{\text{had,LO}}$ corresponds to 0.2%, hardly ever achievable for the absolute measurement of $\sigma(e^+e^- \rightarrow \text{hadrons})$
3. a_μ^{had} calculation from 1st principles (QCD, Lattice).
QCD instanton model (A. Dorokhov, 2003),
Recently from the Lattice: $a_\mu^{\text{had}} = (545 \pm 65) \cdot 10^{-10} \Rightarrow (667 \pm 20) \cdot 10^{-10}$
(C. Aubin and T. Blum, 2005),
attempts to estimate a_μ^{lbl} (M. Hayakawa et al., 2005).

Explaining the Discrepancy: Errors or New Physics?

M.Passera, W.J.Marciano, A.Sirlin, 2008

- The $l\bar{l}$ term is wrong \Rightarrow Should move it by $8-10\sigma$
- If assume errors in $\sigma(s)$ and increase it, $\Delta\alpha_{\text{had}}^{(5)}(M_Z)$ also increases $\Rightarrow M_H^{UB}$ decreases restricting a narrow allowed region $114 \text{ GeV} < M_H < 154 \text{ GeV}$
- Using τ data also increases $\Delta\alpha_{\text{had}}^{(5)}(M_Z)$ and leads to $M_H < 133 \text{ GeV}$
- To bridge Δa_μ , $\sigma(s)$ should be increased by 4% from threshold to infinity. Then $M_H < 70 \text{ GeV}$. If $\sigma(s)$ is increased locally, $M_H < 130 \text{ GeV}$.
- All scenarios look rather unlikely \Rightarrow New Physics?

Conclusions

- BNL success stimulated significant progress of experiment and theory
- QED and EW terms are in good shape
- Improved e^+e^- data \Rightarrow smaller $\delta a_\mu^{\text{had,LO}}$ matching experiment
- τ data could further improve the accuracy but a serious yet unexplained failure of CVC relations for e^+e^- and τ is observed
- Further improvement in $a_\mu^{\text{had,LO}}$ by a factor of 2 will be possible after VEPP-2000, DAΦNE-II, CESRc and $(c - \tau)$ factory + ISR at DAΦNE and B-factories
- Light-by-light term will soon limit the accuracy
- More theory input needed
- $a_\mu^{\text{exp}} - a_\mu^{\text{SM}}$ differs from 0 by $\sim 3 \sigma$ or more \Rightarrow A hint to New Physics?

Back-up Slides

Calculations of $a_{\mu}^{\text{had,LO}}$

Authors	Year	$a_{\mu}^{\text{had,LO}}, 10^{-10}$
C.Bouchiat, L.Michel	1961	$\simeq 648$
M.Gourdin, E. de Rafael	1969	650 ± 50
A.Bramon et al.	1972	680 ± 90
J.Calmet et al.	1977	702 ± 80
T.Kinoshita et al.	1985	707 ± 18
S.Eidelman, F.Jegerlehner	1995	702 ± 15
R.Alemany et al.	1998	701.1 ± 9.4
M.Davier, A.Höcker	1998	692.4 ± 6.2

Light-by-Light Scattering – II

- A. Pivovarov, 2001: $\sim 14.3 \cdot 10^{-10}$
 u, d, s with m_q , heavy quarks with standard masses
- J.F. de Trocóniz, F.J. Ynduráin, 2001: $(9.2 \pm 2.0) \cdot 10^{-10}$
 Constituent quarks + π^0
- A. Dorokhov, 2005: $(10.6 \pm 1.0) \cdot 10^{-10}$
 Instanton liquid model
- J. Erler, G. Toledo Sánchez, 2006: $< 15.9 \cdot 10^{-10}$
 Parton model

Is there a way to consistently relate experimental data on $\gamma\gamma^* \rightarrow X^0$ with $J^{PC} = 0^{-+}, 0^{++}, 1^{++}, 2^{-+}, 2^{++}$ to the corresponding contributions to $a_\mu^{\text{had,LBL}}$?

Branchings of $\tau^- \rightarrow X^- \nu_\tau$ Decay, %

Hadronic State X	Experiment, 2002	CVC Prediction	$\mathcal{B}_{\text{exp}} - \mathcal{B}_{\text{CVC}}$
$\pi^- \pi^0$	25.31 ± 0.18	24.76 ± 0.25	0.55 ± 0.31
$\pi^- 3\pi^0$	1.08 ± 0.10	1.07 ± 0.05	0.01 ± 0.11
$2\pi^- \pi^+ \pi^0$	4.19 ± 0.23	3.84 ± 0.17	0.35 ± 0.29
$\omega \pi^-$	1.94 ± 0.07	1.82 ± 0.07	0.12 ± 0.10
Total	31.59 ± 0.31	30.28 ± 0.34	1.31 ± 0.46


With more accurate data some deviations have been observed.

Corrections to the τ Spectral Functions



- $S_{EW} = 1.0233 \pm 0.0006$
- Real photons, loops
- FSR
- $m_{\pi^\pm} \neq m_{\pi^0}$
(phase space, Γ_ρ)
- $m_{\rho^\pm} \neq m_{\rho^0}$
- $\rho - \omega$ interference
- Radiative decays
($\pi\pi\gamma, \pi(\eta)\gamma, l^+l^-$)
- $m_u \neq m_d$
and 2 class currents

V. Cirigliano, G. Ecker,
H. Neufeld, 2002

M. Davier, S. Eidelman,
A. Höcker, Z. Zhang, 2002




CVC in the 2π Channel. e^+e^- vs. τ




The branching from all groups is systematically higher than the CVC prediction:
 $B_\tau - B_{ee} = (0.94 \pm 0.32)\%$!

2π Spectral Functions in e^+e^- and τ after KLOE and SND



The puzzle remains unsolved \Rightarrow τ data not used


CVC in the 4π Channel. e^+e^- vs. τ

 $2\pi^+2\pi^-$


$$\mathcal{B}(\tau), \% \quad 1.01 \pm 0.08$$

$$\mathcal{B}(\text{CVC}), \% \quad 1.09 \pm 0.08$$

$$\Delta\mathcal{B}, \% \quad -0.08 \pm 0.11$$

 $\pi^+\pi^-2\pi^0$


$$4.54 \pm 0.13$$

$$3.63 \pm 0.21$$


$$+0.91 \pm 0.25$$

Why are e^+e^- and τ Spectral Functions Different?

- Problems with data: underestimated systematics, normalization, rad. corr.
- Problems with SU(2) breaking corrections; Is ChPT reliable? The uncertainty of corrections may be large (K.Maltman, 2005)
- Non (V-A) contribution to e/w interactions (M.Chizhov, 2003) inspired by problems in $\pi^+ \rightarrow e^+ \nu_e \gamma$ (E.Frlez et al., 2003)
- Effect of charged Higgs propagator in τ decay
- $m_{\rho^\pm} > m_{\rho^0}$ by a few MeV (S.Ghozzi and F.Jegerlehner, 2003, M.Davier, 2003). Current experiments indicate equality within a few MeV.

VEPP-2000

Layout of the VEPP-2000 complex



The design luminosity $\mathcal{L} = 10^{32} \text{ cm}^{-2}\text{s}^{-1}$, with $\int L dt \approx 1 - 2 \text{ fb}^{-1}$ during 3-5 years $\Delta a_\mu^{\text{had}} / a_\mu^{\text{had}}$ can be improved by a factor of 2!